Wisconsin AIDS/HIV Surveillance Summary Cases reported 1983 through 2006

EAU CLAIRE COUNTY

| | | Cumulative <u>Cases</u> <u>Percent</u> | | Reported 2001-2005 <u>Cases</u> <u>Percent</u> | | Reported in 2006 <u>Cases</u> <u>Percent</u> | | Presumed alive Cases Percent | |
|--|---------------|---|----|--|---|--|----|---------------------------------|--|
| Total cases | 89 | 100.0% | 19 | 100.0% | 0 | | 51 | 100.0% | |
| Current disease category | | | | | | | | | |
| AIDS | 65 | 73.0% | 10 | 52.6% | 0 | _ | 28 | 54.9% | |
| Non-AIDS | 24 | 27.0% | 9 | 47.4% | 0 | - | 23 | 45.1% | |
| Risk Exposure Categories | | | | | | | | | |
| Men who have sex with men (MSM) | 49 | 55.1% | 7 | 36.8% | 0 | - | 28 | 54.9% | |
| Injecting drug use (IDU) | 6 | 6.7% | 2 | 10.5% | 0 | _ | 4 | 7.8% | |
| MSM and IDU | 11 | 12.4% | 2 | 10.5% | 0 | _ | 5 | 9.8% | |
| Hemophilia/Coagulation disorder | 3 | 3.4% | 0 | 0.0% | 0 | _ | 1 | 2.0% | |
| High-risk heterosexual contact | 11 | 12.4% | 3 | 15.8% | 0 | - | 7 | 13.7% | |
| Transfusion-associated | 2 | 2.2% | 0 | 0.0% | 0 | - | 0 | 0.0% | |
| Mother with/at risk | 0 | 0.0% | 0 | 0.0% | 0 | - | 0 | 0.0% | |
| Undetermined | 7 | 7.9% | 5 | 26.3% | 0 | - | 6 | 11.8% | |
| Gender | | | | | | | | | |
| Female | 9 | 10.1% | 3 | 15.8% | 0 | - | 6 | 11.8% | |
| Male | 80 | 89.9% | 16 | 84.2% | 0 | - | 45 | 88.2% | |
| Unknown | 0 | 0.0% | 0 | 0.0% | 0 | - | 0 | 0.0% | |
| Race/Ethnicity | | | | | | | | | |
| White | 76 | 85.4% | 16 | 84.2% | 0 | _ | 44 | 86.3% | |
| Black | 6 | 6.7% | 1 | 5.3% | 0 | _ | 5 | 9.8% | |
| Hispanic | 4 | 4.5% | 1 | 5.3% | 0 | _ | 1 | 2.0% | |
| Asian/Pacific Islander | 1 | 1.1% | 0 | 0.0% | 0 | _ | 1 | 2.0% | |
| American Indian | 2 | 2.2% | 1 | 5.3% | 0 | _ | 0 | 0.0% | |
| Multi-racial | 0 | 0.0% | 0 | 0.0% | 0 | _ | 0 | 0.0% | |
| Unknown | 0 | 0.0% | 0 | 0.0% | 0 | - | 0 | 0.0% | |
| Age at diagnosis* | | | | | | | | | |
| Under 5 | 0 | 0.0% | 0 | 0.0% | 0 | _ | 0 | 0.0% | |
| 5-14 | 0 | 0.0% | 0 | 0.0% | 0 | _ | 0 | 0.0% | |
| 15-19 | 2 | 2.2% | 1 | 5.3% | 0 | _ | 0 | 0.0% | |
| 20-24 | 10 | 11.2% | 4 | 21.1% | 0 | _ | 0 | 0.0% | |
| 25-44 | 63 | 70.8% | 11 | 57.9% | 0 | - | 29 | 56.9% | |
| 45 years and older | 14 | 15.7% | 3 | 15.8% | 0 | - | 22 | 43.1% | |
| Unknown | 0 | 0.0% | 0 | 0.0% | 0 | - | 0 | 0.0% | |
| * For cases presumed alive, age is the age | on 12/31/2006 | i. | | | | | | | |
| Year of Report | | | | | | | | | |
| Before 1990 | 14 | 15.7% | | | | | | | |
| 1990 | 4 | 4.5% | | | | | | | |
| 1991 | 9 | 10.1% | | | | | | | |
| 1992 | 6 | 6.7% | | | | | | | |
| 1993 | 10 | 11.2% | | | | | | | |
| 1994 | 5 | 5.6% | | - 5 | | | | _ | |
| 1995 | 5 | 5.6% | | | | | | | |
| 1996 | 3 | 3.4% | | | | | | | |
| 1997 | 2 | 2.2% | | | | | | | |
| | 4 | 2.2% 4.5% | | | | | | | |
| 1998 | | | | | | | | | |
| 1999 | 5 | 5.6% | | | | | | | |
| 2000 | 3 | 3.4% | | | | | | 7 | |
| 2001 | 3 | 3.4% | | | | | | | |
| 2002 | 2 | 2.2% | | | | | | | |
| 2003 | 8 | 9.0% | | | | | | II. | |
| 2004 | 2 | 2.2% | | | | | | | |
| 2005 | 4 | 4.5% | | | | | | | |
| 2006 | 0 | 0.0% | | | | | | | |